

Noncommutative Einstein-AdS Gravity in three Dimensions

S. Cacciatori^{1,3*}, D. Klemm^{2,3†}, L. Martucci^{2,3‡} and D. Zanon^{2,3§}

¹ Dipartimento di Matematica dell'Università di Milano,
Via Saldini 50, I-20133 Milano.

² Dipartimento di Fisica dell'Università di Milano,
Via Celoria 16, I-20133 Milano.

³ INFN, Sezione di Milano,
Via Celoria 16, I-20133 Milano.

Abstract

We present a Lorentzian version of three-dimensional noncommutative Einstein-AdS gravity by making use of the Chern-Simons formulation of pure gravity in $2+1$ dimensions. The deformed action contains a real, symmetric metric and a real, antisymmetric tensor that vanishes in the commutative limit. These fields are coupled to two abelian gauge fields. We find that this theory of gravity is invariant under a class of transformations that reduce to standard diffeomorphisms once the noncommutativity parameter is set to zero.

*cacciatori@mi.infn.it

†dietmar.klemm@mi.infn.it

‡luca.martucci@mi.infn.it

§daniela.zanon@mi.infn.it

The field theory limit of open string amplitudes in the presence of a $B_{\mu\nu}$ background gives rise to gauge theories formulated in a noncommutative geometry [1]. The parameter $\theta^{\mu\nu}$ which encodes the noncommutativity properties is essentially given by $B_{\mu\nu}^{-1}$. In a field theory perturbative framework it naturally produces an effective cutoff, $\Delta x^\mu \equiv \theta^{\mu\nu} p_\nu$, that modifies the UV behaviour of the theory [2]. As long as $\theta^{\mu\nu}$ is treated as an external, given background, this amounts to an ad hoc procedure. In order to gain true insights one would like to see noncommutativity/nonlocality entering the game as a dynamical object. One way to make progress in this direction might be to understand how to formulate gravity in a noncommutative geometry.

While noncommutative gauge theories have been studied extensively at the perturbative and nonperturbative level, not much is known about the corresponding formulation of a gravitational theory. One of the main obstacle to overcome is related to the fact that the Moyal product, which implements the noncommutativity, does not maintain reality. In the noncommutative formulation of a gauge theory this does not represent a real problem since the deformed gauge transformations are such that they produce new, real gauge fields. One possible way to preserve reality of the gravitational field is to use explicitly the Seiberg-Witten map [3]. Otherwise it seems that one is forced to complexify the fields [4, 5]. However, complex gravity may be plagued by inconsistencies already at the commutative level [6, 7].

In two and in three dimensions however one can take advantage of the fact that a theory of gravity can be formulated as a gauge theory. In fact since we know how to deform gauge transformations and since the metric does not appear in the volume form, in order to obtain the corresponding noncommutative version of the theory it is sufficient to introduce the $\star$ -product and extend the gauge group appropriately.

In this letter we study noncommutative gravity in three dimensions, while the two-dimensional theory will be treated in a forthcoming paper [8].

It is well-known that in three dimensions pure Einstein gravity can be written as a Chern-Simons theory. More precisely the Einstein-Hilbert action with negative cosmological constant is equivalent to the difference of two $SO(2, 1)$ Chern-Simons actions, modulo boundary terms [9, 10]. This was used in [11]¹ in order to obtain a *Euclidean* version of three-dimensional noncommutative gravity, based on the gauge group $GL(2, \mathbb{C})$. We follow a similar approach and define a three-dimensional *Lorentzian* version of noncommutative gravity. One advantage of our formulation is given by the fact that all the fields are real and the metric is naturally identified. We find a set of transformations which are invariances of the action and reduce to standard diffeomorphisms in the commutative limit. Here we present the main results; more details and insights will be given in [8].

Let us introduce the action

¹For extensions cf. [12].

$$S = \beta \int \text{Tr} \left(A \star dA + \frac{2}{3} A \star A \star A \right) - \beta \int \text{Tr} \left(\tilde{A} \star d\tilde{A} + \frac{2}{3} \tilde{A} \star \tilde{A} \star \tilde{A} \right), \quad (1)$$

where β denotes a dimensionless coupling constant to be specified below. The gauge fields are given by

$$A = \omega + \frac{e}{l} + \frac{i}{2}b, \quad (2)$$

$$\tilde{A} = \omega - \frac{e}{l} + \frac{i}{2}\tilde{b}, \quad (3)$$

with

$$\omega = \omega^a \tau_a, \quad e = e^a \tau_a, \quad b = b\mathbb{I} \quad (4)$$

($a = 0, 1, 2$) taking values in the Lie algebra $\mathfrak{u}(1,1)$, whose generators τ_A ($A = 0, 1, 2, 3$) are given in the appendix. We furthermore defined

$$A \star B \equiv A_\mu \star B_\nu dx^\mu \wedge dx^\nu, \quad (5)$$

where the star denotes the usual Moyal product. Via the isomorphism $so(2,1) \sim su(1,1)$, the algebra $u(1,1)$ is the natural noncommutative generalization of $so(2,1)$. Thus we choose $U(1,1)_\star \times U(1,1)_\star$ as the gauge group for (1)². An element $g \in U(1,1)_\star$ must satisfy the relation

$$g_\star^{-1} = \eta g^\dagger \eta, \quad (6)$$

where $\eta = \text{diag}(-1, 1)$, and $g \star g_\star^{-1} = 1 = g_\star^{-1} \star g$.

All the fields are chosen to be real and this reality is preserved under the gauge transformations

$$\begin{aligned} A &\rightarrow g_\star^{-1} \star A \star g + g_\star^{-1} \star dg, \\ \tilde{A} &\rightarrow \tilde{g}_\star^{-1} \star \tilde{A} \star \tilde{g} + \tilde{g}_\star^{-1} \star d\tilde{g}, \end{aligned}$$

that leave invariant (modulo boundary terms) the action (1).

With the definitions

²In order to formulate a Lorentzian noncommutative gravity with positive cosmological constant, one starts from the $SL(2, \mathbb{C})$ Chern-Simons action, which becomes then $GL(2, \mathbb{C})$ in the noncommutative case, like in Euclidean AdS gravity [11]. A Euclidean noncommutative de Sitter gravity can instead be obtained from a $U(2) \times U(2)$ CS action.

$$\mathcal{R} = d\omega + \omega \stackrel{\star}{\wedge} \omega, \quad (7)$$

$$\mathcal{T} = de + \omega \stackrel{\star}{\wedge} e + e \stackrel{\star}{\wedge} \omega, \quad (8)$$

the action in (1) becomes

$$\begin{aligned} S = & \frac{2\beta}{l} \int \text{Tr} \left(2e \stackrel{\star}{\wedge} \mathcal{R} + \frac{2}{3l^2} e \stackrel{\star}{\wedge} e \stackrel{\star}{\wedge} e \right) \\ & - \frac{\beta}{4} \int \text{Tr} \left(b \stackrel{\star}{\wedge} db + \frac{i}{3} b \stackrel{\star}{\wedge} b \stackrel{\star}{\wedge} b \right) + \frac{\beta}{4} \int \text{Tr} \left(\tilde{b} \stackrel{\star}{\wedge} d\tilde{b} + \frac{i}{3} \tilde{b} \stackrel{\star}{\wedge} \tilde{b} \stackrel{\star}{\wedge} \tilde{b} \right) \\ & + \frac{i\beta}{l} \int \text{Tr} \left\{ \left(\omega \stackrel{\star}{\wedge} e + e \stackrel{\star}{\wedge} \omega \right) \stackrel{\star}{\wedge} (b + \tilde{b}) \right\} \\ & + i\beta \int \text{Tr} \left\{ \left(\omega \stackrel{\star}{\wedge} \omega + \frac{1}{l^2} e \stackrel{\star}{\wedge} e \right) \stackrel{\star}{\wedge} (b - \tilde{b}) \right\}. \end{aligned}$$

Using $\text{Tr}(d\omega) = 0 = \text{Tr}(de)$, this simplifies to

$$\begin{aligned} S = & \frac{2\beta}{l} \int \text{Tr} \left(2e \stackrel{\star}{\wedge} \mathcal{R} + \frac{2}{3l^2} e \stackrel{\star}{\wedge} e \stackrel{\star}{\wedge} e \right) \\ & - \frac{\beta}{4} \int \text{Tr} \left(b \stackrel{\star}{\wedge} db + \frac{i}{3} b \stackrel{\star}{\wedge} b \stackrel{\star}{\wedge} b \right) + \frac{\beta}{4} \int \text{Tr} \left(\tilde{b} \stackrel{\star}{\wedge} d\tilde{b} + \frac{i}{3} \tilde{b} \stackrel{\star}{\wedge} \tilde{b} \stackrel{\star}{\wedge} \tilde{b} \right) \\ & + \frac{i\beta}{l} \int \text{Tr} \left\{ \mathcal{T} \stackrel{\star}{\wedge} (b + \tilde{b}) \right\} + i\beta \int \text{Tr} \left\{ \left(\mathcal{R} + \frac{1}{l^2} e \stackrel{\star}{\wedge} e \right) \stackrel{\star}{\wedge} (b - \tilde{b}) \right\}. \quad (9) \end{aligned}$$

This form of the action suggests to interpret e^a as the dreibein fields and ω , $\mathcal{T}$ and $\mathcal{R}$ as connection, torsion and curvature respectively. In fact one can easily make contact with the usual definitions. We have

$$\begin{aligned} \mathcal{R} &= d\omega + \omega \stackrel{\star}{\wedge} \omega = d\omega^a \tau_a + \omega^a \stackrel{\star}{\wedge} \omega^b \tau_a \tau_b \\ &= R^a \tau_a - \frac{i}{2} \eta_{ab} \omega^a \stackrel{\star}{\wedge} \omega^b \tau_3, \end{aligned} \quad (10)$$

where

$$R^a = d\omega^a - \frac{1}{2} \epsilon^a_{bc} \omega^b \stackrel{\star}{\wedge} \omega^c. \quad (11)$$

If one defines

$$\omega^{ab} = \epsilon^{abc} \omega_c, \quad R^{ab} = \epsilon^{abc} R_c, \quad (12)$$

then

$$R^{ab} = d\omega^{ab} + \frac{1}{2} \left(\omega^a_c \star \omega^{cb} - \omega^b_c \star \omega^{ca} \right). \quad (13)$$

In the commutative limit the above expression reduces to the standard curvature in terms of the spin connection.

Similarly for the torsion we have

$$\mathcal{T} = T^a \tau_a - \frac{i}{2} \left(\omega^a \star e_a + e^a \star \omega_a \right) \tau_3, \quad (14)$$

with

$$T^a = de^a + \frac{1}{2} \left(\omega^a_b \star e^b + e^b \star \omega_b^a \right). \quad (15)$$

Using the above definitions the action in (9) can be written as

$$\begin{aligned} S = & \frac{\beta}{l} \int \epsilon_{abc} \left(e^a \star R^{bc} - \frac{1}{3l^2} e^a \star e^b \star e^c \right) \\ & - \frac{\beta}{2} \int \left(b \star db + \frac{i}{3} b \star b \star b \right) + \frac{\beta}{2} \int \left(\tilde{b} \star d\tilde{b} + \frac{i}{3} \tilde{b} \star \tilde{b} \star \tilde{b} \right) \\ & + \frac{i\beta}{2l} \int \eta_{ab} \left(e^a \star \omega^b + \omega^a \star e^b \right) \star (b + \tilde{b}) \\ & + \frac{i\beta}{2} \int \eta_{ab} \left(\omega^a \star \omega^b + \frac{1}{l^2} e^a \star e^b \right) \star (b - \tilde{b}) \end{aligned} \quad (16)$$

Now the coupling constant β can be identified in terms of the three-dimensional Newton constant as

$$\beta = \frac{l}{16\pi G_N}. \quad (17)$$

The action (16) represents a noncommutative version of Einstein gravity with negative cosmological constant $\Lambda = -1/l^2$, coupled to a pair of $U(1)$ Chern-Simons fields³. In particular $b + \tilde{b}$ couples to the torsion and $b - \tilde{b}$ couples to the curvature. Note that this coupling is a pure noncommutative effect.

From (16) we can compute the equations of motion

$$\frac{\delta S}{\delta e^a} = 0 \Rightarrow R^a + \frac{1}{2l^2} \epsilon^a_{bc} e^b \star e^c - \frac{i}{4} [\omega^a \star (b + \tilde{b}) + (b + \tilde{b}) \star \omega^a]$$

³As before $h \in U(1)$ means that $h^\dagger = h_\star^{-1}$.

$$\begin{aligned}
& -\frac{i}{4l} [e^a \star (b - \tilde{b}) + (b - \tilde{b}) \star e^a] = 0, \\
\frac{\delta S}{\delta \omega^a} = 0 \quad \Rightarrow \quad & T^a - \frac{i}{4} [e^a \star (b + \tilde{b}) + (b + \tilde{b}) \star e^a] \\
& -\frac{il}{4} [\omega^a \star (b - \tilde{b}) + (b - \tilde{b}) \star \omega^a] = 0, \\
\frac{\delta S}{\delta b} = 0 \quad \Rightarrow \quad & F(b) - \frac{i}{2l} \eta_{ab} (\omega^a \star e^b + e^a \star \omega^b) \\
& -\frac{i}{2} \eta_{ab} (\omega^a \star \omega^b + \frac{1}{l^2} e^a \star e^b) = 0, \\
\frac{\delta S}{\delta \tilde{b}} = 0 \quad \Rightarrow \quad & F(\tilde{b}) + \frac{i}{2l} \eta_{ab} (\omega^a \star e^b + e^a \star \omega^b) \\
& -\frac{i}{2} \eta_{ab} (\omega^a \star \omega^b + \frac{1}{l^2} e^a \star e^b) = 0,
\end{aligned} \tag{18}$$

where $F(b) = db + \frac{i}{2} b \star b$, $F(\tilde{b}) = d\tilde{b} + \frac{i}{2} \tilde{b} \star \tilde{b}$. Again one can easily check that in the commutative limit one is left with standard Einstein gravity plus two decoupled $U(1)$ Chern-Simons actions.

A natural definition of a metric follows,

$$G_{\mu\nu} = e_\mu^a \star e_\nu^b \eta_{ab} = g_{\mu\nu} + i b_{\mu\nu}, \tag{19}$$

where

$$g_{\mu\nu} = \frac{1}{2} \eta_{ab} \{e_\mu^a, e_\nu^b\} \tag{20}$$

is real and symmetric, and reduces to the usual expression of the metric in the commutative case, whereas

$$b_{\mu\nu} = -\frac{i}{2} \eta_{ab} [e_\mu^a, e_\nu^b] \tag{21}$$

is real and antisymmetric, and vanishes for $\theta^{\mu\nu} = 0$.

We will now show that the action (1) is invariant under a certain class of deformed transformations that reduce to ordinary diffeomorphisms in the commutative case⁴. We recall that for $\theta^{\mu\nu} = 0$ the variation of a one-form A under an infinitesimal diffeomorphism along an arbitrary vector field v is given by the Lie derivative

$$\mathcal{L}_v A = (di_v + i_v d)A, \tag{22}$$

⁴Cf. also [13].

where i_v is the inner product on differential forms, e. g. $i_v A = v^\rho A_\rho$ for one-forms. It is easy to show that

$$\mathcal{L}_v A = \delta_{i_v A} A + i_v DA, \quad (23)$$

where $DA = dA + A \wedge A$ and $\delta_{i_v A} A$ denotes the variation of A under an infinitesimal gauge transformation with parameter $i_v A$. A natural generalization of (23) to the non-commutative case is

$$\Delta_v^* := \delta_{i_v^* A} A + i_v^* DA, \quad (24)$$

where, for p -forms ω^p ,

$$i_v^* \omega^p := \frac{1}{2(p-1)!} [v^\rho \star \omega_{\rho\mu_1 \dots \mu_{p-1}}^p + \omega_{\rho\mu_1 \dots \mu_{p-1}}^p \star v^\rho] dx^{\mu_1} \wedge \dots \wedge dx^{\mu_{p-1}}, \quad (25)$$

and $DA = dA + A \overset{\star}{\wedge} A =: F$. A more detailed discussion of the deformed transformations (24) will be presented in [8]. It is clear by construction that the transformations (24) reduce to ordinary diffeomorphisms in the commutative limit.

From gauge invariance of the action (1) and from (24) it follows that it is sufficient to prove that the action is invariant (modulo boundary terms) under the transformations

$$\begin{aligned} \delta'_v A &= i_v^* DA = i_v^* F, \\ \delta'_v \tilde{A} &= i_v^* D\tilde{A} = i_v^* \tilde{F}, \end{aligned}$$

then also invariance under (24) holds. It is straightforward to show that

$$\begin{aligned} \delta'_v S &= 2\beta \int \text{Tr} \left(F \overset{\star}{\wedge} i_v^* F \right) - 2\beta \int \text{Tr} \left(\tilde{F} \overset{\star}{\wedge} i_v^* \tilde{F} \right) \\ &= -2\beta \int \text{Tr} \left(i_v^* F \overset{\star}{\wedge} F \right) + 2\beta \int \text{Tr} \left(i_v^* \tilde{F} \overset{\star}{\wedge} \tilde{F} \right) = 0. \end{aligned} \quad (26)$$

We are now interested in how $U(1,1) \times U(1,1)$ gauge transformations act on the fields ω , e , b , $\tilde{b}$. Let

$$\lambda = \lambda^A \tau_A = \lambda^a \tau_a + \chi \tau_3 = (\xi^a + \alpha^a) \tau_a + \chi \tau_3, \quad (27)$$

$$\tilde{\lambda} = \tilde{\lambda}^A \tau_A = \tilde{\lambda}^a \tau_a + \tilde{\chi} \tau_3 = (\xi^a - \alpha^a) \tau_a + \tilde{\chi} \tau_3 \quad (28)$$

be the generators of the infinitesimal gauge transformations, written in terms of the parameters ξ^a , α^a , and $(\chi, \tilde{\chi})$, which refer to rotations, translations and $U(1) \times U(1)$ gauge transformations respectively. Then

$$\begin{aligned}\delta_\lambda A^a &= d\lambda^a - \frac{1}{2}\epsilon^a_{bc} (A^b \star \lambda^c - \lambda^b \star A^c) \\ &\quad + \frac{i}{2} (A^a \star \chi - \chi \star A^a) + \frac{i}{2} (b \star \lambda^a - \lambda^a \star b) ,\end{aligned}\tag{29}$$

$$\delta_\lambda b = d\chi - \frac{i}{2}\eta_{ab} (A^a \star \lambda^b - \lambda^a \star A^b) + \frac{i}{2} (b \star \chi - \chi \star b) ,\tag{30}$$

with analogous relations for $\tilde{A}^a, \tilde{b}$. Furthermore we have

$$\tilde{\delta}_{\tilde{\lambda}} A = \delta_\lambda \tilde{A} = 0 .\tag{31}$$

In what follows it will be convenient to introduce the definitions $\hat{\delta}_{(\lambda, \tilde{\lambda})} := \delta_\lambda + \tilde{\delta}_{\tilde{\lambda}}$, with λ and $\tilde{\lambda}$ as in (27-28), and the combined fields $b := b_1 + b_2$ and $\tilde{b} := b_1 - b_2$.

Let us now consider separately rotations, translations and $U(1) \times U(1)$ gauge transformations.

Rotations:

Rotations are given by setting $\alpha^a = \chi = \tilde{\chi} = 0$, i. e. , by taking transformations with parameters

$$\lambda = \xi = \xi^a \tau_a , \quad \tilde{\lambda} = \xi = \xi^a \tau_a .\tag{32}$$

Defining $\xi_{ab} = \epsilon_{abc} \xi^c$, we obtain for the variation of the spin connection $\omega^{ab} = \epsilon^{abc} \omega_c$

$$\hat{\delta}_{(\xi, \xi)} \omega^{ab} = d\xi^{ab} + \omega^{[a}_c \star \xi^{c|b]} - \xi^{[a}_c \star \omega^{c|b]} + \frac{i}{2} (b_1 \star \xi^{ab} - \xi^{ab} \star b_1) ,\tag{33}$$

and for the dreibein fields

$$\hat{\delta}_{(\xi, \xi)} e^a = -\frac{1}{2} (\xi^a_b \star e^b + e^b \star \xi^a_b) + \frac{i l}{2} (b_2 \star \xi^a - \xi^a \star b_2) .\tag{34}$$

Finally the fields b_1, b_2 transform as

$$\hat{\delta}_{(\xi, \xi)} b_1 = \frac{i}{4} (\xi^a_b \star \omega^b_a - \omega^a_b \star \xi^b_a) ,\tag{35}$$

$$\hat{\delta}_{(\xi, \xi)} b_2 = -\frac{i}{4l} \epsilon_{abc} (\xi^{ab} \star e^c - e^c \star \xi^{ab}) .\tag{36}$$

Translations:

Translations are generated by the fields

$$\lambda^a = \alpha^a, \quad \tilde{\lambda}^a = -\alpha^a, \quad (37)$$

which applied to the spin connection yield

$$\hat{\delta}_{(\alpha, -\alpha)} \omega^{ab} = \frac{1}{l} (e^{[a} \star \alpha^{b]} - \alpha^{[a} \star e^{b]}) + \frac{i}{2} \epsilon^{abc} (b_2 \star \alpha_c - \alpha_c \star b_2). \quad (38)$$

The dreibein transforms according to

$$\hat{\delta}_{(\alpha, -\alpha)} e^a = l d\alpha^a + \frac{l}{2} (\omega^a_b \star \alpha^b - \alpha^b \star \omega_b^a) + \frac{il}{2} (b_1 \star \alpha^a - \alpha^a \star b_1), \quad (39)$$

whereas the variations of the fields b_1, b_2 are given by

$$\hat{\delta}_{(\alpha, -\alpha)} b_1 = \frac{i}{2l} (\alpha_a \star e^a - e^a \star \alpha_a), \quad (40)$$

$$\hat{\delta}_{(\alpha, -\alpha)} b_2 = \frac{i}{4} \epsilon^{abc} (\omega_{ab} \star \alpha_c - \alpha_c \star \omega_{ab}). \quad (41)$$

$U(1) \times U(1)$ transformations:

$U(1) \times U(1)$ transformations are given by the parameters

$$\lambda = \chi \tau_3, \quad \tilde{\lambda} = \tilde{\chi} \tau_3. \quad (42)$$

If we introduce the combinations

$$\chi = \chi_1 + \chi_2, \quad \tilde{\chi} = \chi_1 - \chi_2, \quad (43)$$

the spin connection and the dreibein transform as

$$\begin{aligned} \hat{\delta}_{(\chi, \tilde{\chi})} \omega^{ab} &= \frac{i}{2} (\omega^{ab} \star \chi_1 - \chi_1 \star \omega^{ab}) + \frac{i}{2l} \epsilon^{abc} (e_c \star \chi_2 - \chi_2 \star e_c), \\ \hat{\delta}_{(\chi, \tilde{\chi})} e^a &= \frac{i}{2} (e^a \star \chi_1 - \chi_1 \star e^a) - \frac{il}{4} \epsilon^{abc} (\omega_{bc} \star \chi_2 - \chi_2 \star \omega_{bc}), \end{aligned} \quad (44)$$

whereas the variations of the $U(1)$ Chern-Simons fields read

$$\hat{\delta}_{(\chi, \tilde{\chi})} b = d\chi + \frac{i}{2} (b \star \chi - \chi \star b), \quad (45)$$

$$\hat{\delta}_{(\chi, \tilde{\chi})} \tilde{b} = d\tilde{\chi} + \frac{i}{2} (\tilde{b} \star \tilde{\chi} - \tilde{\chi} \star \tilde{b}). \quad (46)$$

The main goal reached in our work is the formulation of a noncommutative version of Lorentzian Einstein-AdS gravity in three dimensions in terms of real fields. This has been achieved in two steps: first we have used the Chern-Simons gauge theory formulation of gravity and the fact that gauge transformations maintain the reality properties of the fields even in the noncommutative case. Secondly we have defined the gravity fields in such a way that the identification of a real, symmetric metric and a real antisymmetric tensor naturally arise. We have found that this deformed Einstein action possesses a class of invariances that reduce to standard diffeomorphisms in the commutative limit. We have computed how all the relevant fields transform under rotations, translations and $U(1) \times U(1)$ gauge transformations. An open question is still how to quantize the theory and understand which is the dynamical role played by the antisymmetric $b_{\mu\nu}$ field. In order to complete this program one would need to obtain a noncommutative version of general coordinate transformations. Our deformed transformations (which are active) may be related to a subset of them, i. e. , to the ones that maintain the Poisson structure introduced by $\theta^{\mu\nu}$. Another interesting issue to address is how to couple gravity to matter fields.

Acknowledgements

This work was partially supported by INFN, MURST and by the European Commission RTN program HPRN-CT-2000-00131, in which S. C. , D. K. , L. M. and D. Z. are associated to the University of Torino.

A Conventions

An element M of the Lie algebra $u(1,1)$ satisfies

$$M^i_j = -\eta_{jk} \bar{M}^k_l \eta^{li}, \quad (1)$$

where a bar denotes complex conjugation, and $(\eta_{ij}) = \text{diag}(-1, 1)$. We choose as $u(1,1)$ generators

$$\begin{aligned} \tau_0 &= \frac{1}{2} \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, & \tau_1 &= \frac{1}{2} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \\ \tau_2 &= \frac{1}{2} \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, & \tau_3 &= \frac{1}{2} \begin{pmatrix} i & 0 \\ 0 & i \end{pmatrix}. \end{aligned} \quad (2)$$

They are normalized according to

$$\text{Tr}(\tau_A \tau_B) = \frac{1}{2} \eta_{AB}, \quad (3)$$

where $(\eta_{AB}) = \text{diag}(-1, 1, 1, -1)$ is the inner product on the Lie algebra. The generators (2) satisfy the relation (1). Further, if a, b, c assume the values 0, 1 and 2, then the following relations hold:

$$[\tau_a, \tau_b] = -\epsilon_{abc} \tau^c, \quad (4)$$

$$[\tau_a, \tau_3] = 0, \quad (5)$$

$$\tau_a \tau_b = -\frac{1}{2} \epsilon_{abc} \tau^c - \frac{i}{2} \eta_{ab} \tau_3, \quad (6)$$

$$\text{Tr}(\tau_a \tau_b \tau_c) = -\frac{1}{4} \epsilon_{abc}, \quad (7)$$

$$\text{Tr}(\tau_a \tau_b \tau_3) = \frac{i}{4} \eta_{ab}, \quad (8)$$

where $(\eta_{ab}) = \text{diag}(-1, 1, 1)$.

Furthermore we defined $\epsilon_{012} = 1$.

References

- [1] N. Seiberg and E. Witten, JHEP **9909** (1999) 032 [arXiv:hep-th/9908142].
- [2] S. Minwalla, M. Van Raamsdonk and N. Seiberg, JHEP **0002** (2000) 020 [arXiv:hep-th/9912072].
- [3] A. H. Chamseddine, Phys. Lett. B **504** (2001) 33 [arXiv:hep-th/0009153].
- [4] A. H. Chamseddine, Commun. Math. Phys. **218** (2001) 283 [arXiv:hep-th/0005222].
- [5] A. H. Chamseddine, Int. J. Mod. Phys. A **16** (2001) 759 [arXiv:hep-th/0010268].
- [6] T. Damour, S. Deser and J. G. McCarthy, Phys. Rev. D **47** (1993) 1541 [arXiv:gr-qc/9207003].
- [7] J. W. Moffat, J. Math. Phys. **36** (1995) 3722 [Erratum-ibid. **36** (1995) 7128];
Phys. Lett. B **491** (2000) 345 [arXiv:hep-th/0007181].
- [8] S. Cacciatori, A. H. Chamseddine, D. Klemm, L. Martucci, W. A. Sabra and D. Zanon, to appear.
- [9] A. Achúcarro and P. K. Townsend, Phys. Lett. B **180** (1986) 89.
- [10] E. Witten, Nucl. Phys. B **311** (1988) 46.
- [11] M. Bañados, O. Chandia, N. Grandi, F. A. Schaposnik and G. A. Silva, Phys. Rev. D **64** (2001) 084012 [arXiv:hep-th/0104264].
- [12] V. P. Nair, arXiv:hep-th/0112114.
- [13] R. Jackiw and S. Y. Pi, arXiv:hep-th/0111122.